



## Head & neck primary cutaneous melanoma patients stratified by CP-GEP (Merlin Assay): risk of nodal metastasis and long-term survival outcome

M.R. Pittelkow<sup>a</sup>, W. Yu<sup>b</sup>, T.J. Hieken<sup>c</sup>, R. Wever<sup>d</sup>, F. Tjien-Fooh<sup>d</sup>, J. Dwarkasing<sup>d</sup>, A. Meves<sup>e</sup>

a Department of Dermatology, Mayo Clinic, Scottsdale, Arizona, USA

b Department of Dermatology, Oregon Health and Science University, Portland, Oregon, USA

c Department of Surgery, Mayo Clinic, Rochester, MN, USA

d SkylineDx B.V., 3062 ME Rotterdam, The Netherlands

e Department of Dermatology, Mayo Clinic, Rochester, Minnesota, USA

Abstract at SSO 2024

## *Introduction*

Primary cutaneous melanoma (CM) of the head and neck often yields false-negative results from sentinel lymph node biopsies (SLNB). Consequently, these melanomas exhibit higher recurrence rates than those on other body parts. In this study, we investigate the capability of CP-GEP to stratify patients' risk for nodal metastasis and evaluate their long-term survival outcomes.

## *Methods*

Patients with primary CM of the head and neck who underwent SLNB between 2004 and 2021 at two US centers were included in this study. CP-GEP integrates Breslow thickness, the patient's age at diagnosis, and the expression of eight genes from the primary CM tissue. The CP-GEP model produces a binary output: High Risk or Low Risk.

## *Results*

Out of 250 head and neck CM patients included in the analysis, there was a 14.0% SLNB positivity rate. CP-GEP classified 147 patients (58.8%) as High Risk. These patients exhibited a higher SLNB positivity rate of 22.4% and accounted for 14 of the 19 melanoma-specific deaths (73.7%). Meanwhile, CP-GEP categorized 103 patients as Low Risk, leading to a 41.2% reduction in SLNB with a Negative Predictive Value (NPV) of 98.1%. The five-year Melanoma-Specific Survival (MSS) rates were 82.4% for High Risk patients and 95.5% for Low Risk patients.

## *Conclusion*

CP-GEP has the potential to categorize primary CM patients of the head and neck based on their risk for nodal metastasis and long-term survival outcomes. Notably, CP-GEP can help identify patients with tumors in surgically challenging locations who potentially may avoid SLNB surgery, while more accurately pinpointing those truly at risk for melanoma-specific death.

---

